arXiv:1704.04018v1 [math.RT] 13 Apr 2017

The Fourier transform on the group $\mathrm{GL}_2(\mathbb{R})$ and the action of the overalgebra $\mathfrak{gl}_4$

Yu.A.Neretin[1]

We define a kind of 'operational calculus' for $\mathrm{GL}_2(\mathbb{R})$. Namely, the group $\mathrm{GL}_2(\mathbb{R})$ can be regarded as an open dense chart in the Grassmannian of 2-dimensional subspaces in $\mathbb{R}^4$. Therefore the group $\mathrm{GL}_4(\mathbb{R})$ acts in L^2 on $\mathrm{GL}_2(\mathbb{R})$. We transfer the corresponding action of the Lie algebra $\mathfrak{gl}_4$ to the Plancherel decomposition of $\mathrm{GL}_2(\mathbb{R})$, the algebra acts by differential-difference operators with shifts in an imaginary direction. We also write similar formulas for the action of $\mathfrak{gl}_4 \oplus \mathfrak{gl}_4$ in the Plancherel decomposition of $\mathrm{GL}_2(\mathbb{C})$.

1 The statement of the paper

1.1. The group $\mathrm{GL}_(\mathbb{R})$. Let $\mathrm{GL}_n(\mathbb{R})$ be the group of invertible real matrices of order n. We denote elements of $\mathrm{GL}_2(\mathbb{R})$ by

$$X = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}.$$

The Haar measure on $\mathrm{GL}_2(\mathbb{R})$ is given by

$$(\det X)^{-2}\, dX = (\det X)^{-2}\, dx_{11}\, dx_{12}\; dx_{21}\, dx_{22}.$$

Recall some basic facts on representations of the group $\mathrm{GL}_2(\mathbb{R})$, for systematic exposition of the representation theory of $\mathrm{SL}_2(\mathbb{R})$, see, e.g., [5], [7]. Denote by $\mathbb{R}^\times$ the multiplicative group of $\mathbb{R}$. Let $\mu \in \mathbb{C}$ and $\varepsilon \in \mathbb{Z}/2\mathbb{Z}$. We define the function $x^{\mu /\!\!/ \varepsilon}$ on $\mathbb{R}^\times$ by

$$x^{\mu /\!\!/ \varepsilon} := |x|^\mu \operatorname{sgn}(x)^\varepsilon.$$

Denote by Λ the set of all collections

$$(\mu_1, \varepsilon_1; \mu_2, \varepsilon_2),$$

i.e.,

$$\Lambda \simeq \mathbb{C} \times \mathbb{Z}_2 \times \mathbb{C} \times \mathbb{Z}_2.$$

For each element of Λ we define a representation $T_{\mu,\varepsilon}$ of $\mathrm{GL}_2(\mathbb{R})$ in the space of functions on $\mathbb{R}$ by

$$T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} \varphi(t) =$$
$$= \varphi\Big(\frac{x_{12}+tx_{22}}{x_{11}+tx_{21}}\Big) \cdot (x_{11}+tx_{21})^{-1+\mu_1-\mu_2 /\!\!/ \varepsilon_1-\varepsilon_2} \det \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}^{1/2+\mu_2 /\!\!/ \varepsilon_2}. \tag{1.1}$$

[1]Supported by the grant FWF, Project P28421.

A space of functions can be specified in various ways, it is convenient to consider the space $C^\infty_{\mu_1-\mu_2,\varepsilon_1-\varepsilon_2}$ of C^∞-functions on $\mathbb{R}$ such that[2]

$$\varphi(-1/t)(-t)^{-1+\mu_1-\mu_2/\!\!/\varepsilon_1-\varepsilon_2}$$

also is C^∞-smooth.

Thus, for any fixed $X \in \mathrm{GL}_2(\mathbb{R})$ we get an operator-valued function $(\mu_1, \varepsilon_1; \mu_2, \varepsilon_2) \mapsto T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}$ on Λ holomorphic in the variables μ_1, μ_2.

Generators of the Lie algebra $\mathfrak{gl}_2(\mathbb{R})$ act by formulas

$$L_{11} = -t\frac{d}{dt} + (-1/2 + \mu_1), \qquad L_{12} = \frac{d}{dt}, \tag{1.2}$$

$$L_{21} = -t^2\frac{d}{dt} + t(-1 + \mu_1 - \mu_2), \qquad L_{22} = t\frac{d}{dt} + (1/2 + \mu_2). \tag{1.3}$$

The expressions for the generators do not depend on ε_1, ε_2. However the space of the representation depends on $\varepsilon_1 - \varepsilon_2$.

Consider integral operators

$$A_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} : C^\infty_{\mu_1-\mu_2,\varepsilon_1-\varepsilon_2} \to C^\infty_{\mu_2-\mu_1,\varepsilon_1-\varepsilon_2}$$

defined by

$$A_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} f(t) := \int_{\mathbb{R}} (t-s)^{-1-\mu_1+\mu_2/\!\!/\varepsilon_1-\varepsilon_2} f(s)\, ds.$$

The integral is convergent if $\mathrm{Re}(\mu_1 - \mu_2) < 0$ and determines a function holomorphic in μ_1, μ_2. As usual (see, e.g., [4], §I.3), the integral admits a meromorphic continuation to the whole plane $(\mu_1, \mu_2) \in \mathbb{C}^2$ with poles at $\mu_1 - \mu_2 = 0, 1, 2, \ldots$. The operators $A_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}$ are intertwinig,

$$A_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} = T_{\mu_2,\varepsilon_2;\mu_1,\varepsilon_1} A_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2} \tag{1.4}$$

(parameters (μ_1, ε_1) and (μ_2, ε_2) of T are transposed). If $\mu_1 - \mu_2 \neq \pm 1, \pm 2, \ldots$, then representations $T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}$ are irreducible and operators $T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}$ are invertible.

Representations $T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}$ form a so-called (nonunitary) *principal series of representations.* Recall description of some unitary irreducible representations of $\mathrm{GL}_2(\mathbb{R})$.

Unitary principal series. If μ_1, $\mu_2 \in i\mathbb{R}$, then the representation is unitary in $L^2(\mathbb{R})$. Denote by $\Lambda_{\mathrm{principal}} \subset \Lambda$ the subset consisting of tuples $(is_1, \varepsilon_1; is_2, \varepsilon_2)$ with $s_1 \geqslant s_2$.

Discrete series. Notice that the group $\mathrm{GL}_2(\mathbb{R})$ acts on the Riemann sphere $\overline{C} = \mathbb{C} \cup \infty$ by linear-fractional transformations

$$X : z \mapsto \frac{x_{12} + zx_{22}}{x_{11} + zx_{21}}, \tag{1.5}$$

[2]This condition means that functions φ are smooth as sections of line bundles on the projective line $\mathbb{R} \cup \infty$.

these transformations preserve the real projective line $\mathbb{R} \cup \infty$ and therefore preserves its complement $\mathbb{C} \setminus \mathbb{R}$. If $\det X > 0$, then (1.5) leave upper and lower half-planes invariant, if $\det X < 0$, then (1.5) permutes the half-planes.

Let $n = 1, 2, 3, \ldots$. Consider the space H_n of holomorphic functions φ on $\mathbb{C} \setminus \mathbb{R}$ satisfying

$$\int_{\mathbb{C}\setminus\mathbb{R}} |\varphi(z)|^2 |\operatorname{Im} z|^{-n-1} \, d\operatorname{Re} z \, d\operatorname{Im} z < \infty.$$

In fact, φ is a pair of holomorphic functions φ_+ and φ_- determined on half-planes $\operatorname{Im} z > 0$ and $\operatorname{Im} z < 0$. These functions have boundary values on $\mathbb{R}$ in distributional sense (we omit a precise discussion since it is not necessary for our purposes). The space H_n is a Hilbert space with respect to the inner product

$$\langle \varphi_1, \varphi_2 \rangle = \int_{\mathbb{C}\setminus\mathbb{R}} \varphi_1(z) \overline{\varphi_2(z)} \, |\operatorname{Im} z|^{-n-1} \, d\operatorname{Re} z \, d\operatorname{Im} z.$$

For $s \in \mathbb{R}$, $\delta \in \mathbb{Z}_2$ we define a unitary representation $D_{n,s}$ of $\mathrm{GL}_2(\mathbb{R})$ in H_n by

$$D_{n,s} \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} \varphi(z) = \varphi\Big(\frac{x_{12} + z x_{22}}{x_{11} + z x_{21}}\Big) (x_{11} + z x_{21})^{-1-n} \times \\ \times \det \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}^{1/2+n/2+is/\!\!/\delta}.$$

In fact, we have operators (1.1) for

$$\mu_1 = -n/2 - is, \, \varepsilon_1 = -1 + n + \delta; \;\; \mu_2 = n/2 + is, \, \varepsilon_2 = \delta$$

restricted to the subspace generated by boundary values of functions f_+ and f_-.

We denote by $\Lambda_{\mathrm{discrete}}$ the set of all parameters (n, is) of discrete series.

1.2. The Fourier transform. Let φ be contained in the space $C_0^\infty(\mathrm{GL}_2(\mathbb{R}))$ of compactly supported function on $\mathrm{GL}_2(\mathbb{R})$. We consider a function that sent each $\widetilde{\mu} = (\mu_1, \varepsilon_1; \mu_2, \varepsilon_2)$ to an operator in $C^\infty_{\mu_1-\mu_2, \varepsilon_1-\varepsilon_2}$ given by

$$T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}(F) = \int_{\mathrm{GL}_2(\mathbb{R})} F(X) \, T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}(X) \, \frac{dX}{\det(X)^2}.$$

Next, we define a subset $\Lambda_{\mathrm{tempered}} \subset \Lambda$ by

$$\Lambda_{\mathrm{tempered}} := \Lambda_{\mathrm{principal}} \cup \Lambda_{\mathrm{discrete}}.$$

Let us define *Plancherel measure* $d\mathcal{P}(\widetilde{\mu})$ on $\Lambda_{\mathrm{tempered}}$. On the piece $\Lambda_{\mathrm{principal}}$ it is given by

$$d\mathcal{P}(is_1, 0; is_2, \varepsilon_2) = \frac{1}{16\pi^3} (s_1 - s_2) \tanh \pi(s_1 - s_2)/2 \, ds_1 \, ds_2;$$
$$d\mathcal{P}(is_1, 1; is_2, \varepsilon_2) = \frac{1}{16\pi^3} (s_1 - s_2) \coth \pi(s_1 - s_2)/2 \, ds_1 \, ds_2.$$

On n-th piece of $\Lambda_{\text{discrete}}$ it is given by

$$d\mathcal{P} = \frac{n}{8\pi^3} ds.$$

Consider the space $\mathcal{L}^2$ of functions Q on $\Lambda_{\text{tempered}}$ taking values in the space of Hilbert–Schmidt operators in the corresponding Hilbert spaces and satisfying the condition

$$\int_{\Lambda_{\text{tempered}}} \operatorname{tr}\Big(Q(\widetilde{\mu})Q^*(\widetilde{\mu})\Big)\, d\mathcal{P}(\widetilde{\mu}) < \infty.$$

This is a Hilbert space with respect to the inner product

$$\langle Q_1, Q_2\rangle_{\mathcal{L}^2} := \int_{\Lambda_{\text{tempered}}} \operatorname{tr}\Big(Q_1(\widetilde{\mu})Q_2^*(\widetilde{\mu})\Big)\, d\mathcal{P}(\widetilde{\mu}).$$

According the Plancherel theorem, *for any* F_1, $F_2 \in C_0^\infty(\mathrm{GL}_2(\mathbb{R}))$ *we have*

$$\langle F_1, F_2\rangle_{L^2(\mathrm{GL}_2(\mathbb{R}))} = \langle T(F_1), T(F_2)\rangle_{\mathcal{L}^2(\Lambda_{\text{tempered}})}.$$

Moreover, the map $F \to T(F)$ *extends to a unitary operator* $L^2(\mathrm{GL}_2(\mathbb{R})) \to \mathcal{L}$.

1.3. Overgroup. Let $\mathrm{Mat}_2(\mathbb{R})$ be the space of all real matrices of order 2. By $\mathrm{Gr}_{4,2}(\mathbb{R})$ we denote the Grassmannian of 2-dimensional subspaces in $\mathbb{R}^2 \oplus \mathbb{R}^2$. For any operator $\mathbb{R}^2 \to \mathbb{R}^2$ its graph is an element $\mathrm{Gr}_{4,2}(\mathbb{R})$. The set $\mathrm{Mat}_2(\mathbb{R})$ of such operators is an open dense chart in $\mathrm{Gr}_{4,2}(\mathbb{R})$.

The group $\mathrm{GL}_4(\mathbb{R})$ acts in a natural way in $\mathbb{R}^4$ and therefore on the Grassmannian. In the chart $\mathrm{Mat}_2(\mathbb{R})$ the action is given by the formula (see, e.g., [18], Theorem 2.3.2)

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} : X \mapsto (A + XC)^{-1}(B + XD),$$

where $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is an element of $\mathrm{GL}_4(\mathbb{R})$ written as a block matrix of size $2+2$. The Jacobian of this transformation is (see, e.g., [18], Theorem 2.3.2)

$$\det(A + XC)^{-4} \det \begin{pmatrix} A & B \\ C & D \end{pmatrix}^2.$$

For $\sigma \in i\mathbb{R}$ we define a unitary representation of $\mathrm{GL}_4(\mathbb{R})$ in $L^2(\mathrm{Mat}_2(\mathbb{R}))$ by

$$R_\sigma \begin{pmatrix} A & B \\ C & D \end{pmatrix} F(X) = F\big((A + XC)^{-1}(B + XD)\big) \times \\ \times\ |\det(A + XC)|^{-2+2\sigma} \left|\det \begin{pmatrix} A & B \\ C & D \end{pmatrix}\right|^{1-\sigma}$$

(these representations are contained in degenerate principal series).

The group $\mathrm{GL}_2(\mathbb{R})$ is an open dense subset in $\mathrm{Mat}_2(\mathbb{R})$. Therefore, we can identify the spaces L^2 on $\mathrm{GL}_2(\mathbb{R})$ and $\mathrm{Mat}_2(\mathbb{R})$. For this we consider a unitary operator

$$J:\, L^2\big(\mathrm{Mat}_2(\mathbb{R})\big) \to L^2\big(\mathrm{GL}_2(\mathbb{R})\big)$$

given by

$$J_\sigma F(X) = F(X)\cdot |\det X|^{1-\sigma}$$

This determines a unitary representation $U_\sigma := J_\sigma R_\sigma J_\sigma^{-1}$ of $\mathrm{GL}_4(\mathbb{R})$ in $L^2\big(\mathrm{GL}_2(\mathbb{R})\big)$, the explicit expression is

$$U_\sigma \begin{pmatrix} A & B \\ C & D \end{pmatrix} F(X) = F\big((A+XC)^{-1}(B+XD)\big)\times \\ \times \left|\frac{\det(A+XC)\det(B+XD)}{\det X \det\begin{pmatrix} A & B \\ C & D \end{pmatrix}}\right|^{-1+\sigma}. \quad (1.6)$$

Consider a subgroup $\mathrm{GL}_2(\mathbb{R})\times \mathrm{GL}_2(\mathbb{R}) \subset \mathrm{GL}_4(\mathbb{R})$ consisting of matrices $\begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix}$. For this subgroup we get the usual left-right action in $L^2\big(\mathrm{GL}_2(\mathbb{R})\big)$,

$$U_\sigma \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} F(X) = F(A^{-1}XD).$$

Formula (1.6) extends this formula to the whole group $\mathrm{GL}_4(\mathbb{R})$. The Lie algebra $\mathfrak{gl}_4$ acts in the space of functions on $\mathrm{Mat}_2(\mathbb{R})$ by first order differential operators, which can be easily written; a list of formulas for all generators e_{kl}, where k, $l = 1, 2, 3, 4$, is given below in Subs. (2.5). We restrict this action to the space of smooth compactly supported functions on $\mathrm{GL}_2(\mathbb{R})$. Notice that the operators $i\cdot e_{kl}$ are symmetric on this domain, but some of them are not essentially self-adjoint.

Our purpose is to write explicitly the images E_{kl} of operators e_{kl} under the Fourier transform.

1.4. Formulas. Operators $T_{\mu_1,\varepsilon_1;\mu_2;\varepsilon_2}(F)$ have the form

$$T_{\mu_1,\varepsilon_1;\mu_2;\varepsilon_2}(F)\varphi(t) = \int_{-\infty}^{\infty} K(t,s|\mu_1,\varepsilon_1;\mu_2,\varepsilon_2)\,\varphi(t)\,dt. \quad (1.7)$$

Recall that functions K are holomorphic in μ_1, μ_2.

We wish to write operators E_{kl} on kernels K. The complete list is contained below in Subsect. 2.5, here we present two basic expressions.

The algebra $\mathfrak{gl}_4(\mathbb{R})$ can be decomposed as a linear space into a direct sum of four subalgebras $\mathfrak{a}$, $\mathfrak{b}$, $\mathfrak{c}$, $\mathfrak{d}$ consisting of matrices of the form

$$\begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}, \qquad \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix}, \qquad \begin{pmatrix} 0 & 0 \\ * & 0 \end{pmatrix}, \qquad \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix}$$

The subalgebras $\mathfrak{a}$ and $\mathfrak{d}$ are isomorphic to $\mathfrak{gl}_2(\mathbb{R})$, subalgebras $\mathfrak{b}$ and $\mathfrak{c}$ are Abelian.

Formulas for the action of $\mathfrak{a}$ and $\mathfrak{d}$ immediately follow from the definition of the Fourier transform. To obtain formulas for the whole $\mathfrak{gl}_4$, it is sufficient to write expressions for one generator of $\mathfrak{b}$, say E_{14}, and one generator of $\mathfrak{c}$, say E_{32}. After this other generators can be obtained by evaluation of commutators.

We define shift operators V_1^+, V_1^-, V_2^+, V_2^- by

$$V_1^{\pm} K(t,s|\mu_1,\varepsilon_1;\mu_2,\varepsilon_2) = K(t,s|\mu_1 \pm 1,\varepsilon_1+1;\mu_2,\varepsilon_2);$$
$$V_2^{\pm} K(t,s|\mu_1,\varepsilon_1;\mu_2,\varepsilon_2) = K(t,s|\mu_1,\varepsilon_1;\mu_2 \pm 1,\varepsilon_2+1).$$

In this notation,

$$E_{14} = \frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\,V_1^- + \frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\,V_2^-;$$

$$E_{32} = \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\,V_1^+ + \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\,V_2^+.$$

1.5. Remarks on a general problem. In [17] the author formulated the following question: *Assume that we know the explicit Plancherel formula for the restriction of a unitary representation ρ of a group G to a subgroup H. Is it possible to write the action of the Lie algebra of G in the direct integral of representations of H?*

Now it seems that an answer to this question is affirmative.

The initial paper [17] contains a solution for a tensor product[3] of a highest and lowest weight representations of $\mathrm{SL}_2(\mathbb{R})$. In this case the overalgebra acts by differential-difference operators in the space $L^2(S^1 \times \mathbb{R}_+)$ having the form

$$Lf(\varphi,s) = D_1 f(\varphi,s+i) + D_2 f(\varphi,s) + D_3 f(\varphi,s-i),$$

where D_j are second order differential operators in the variable φ.

In [10]–[13] Molchanov solved several rank 1 problems of this type, expressions are similar, but there appear differential operators of order 4. In [20] the action of the overalgebra in restrictions from $\mathrm{GL}_{n+1}(\mathbb{C})$ to $\mathrm{GL}_n(\mathbb{C})$, in this case differential operators have order n. In all the cases examined by now shift operators in the imaginary direction are present.

In the present paper, we write the action of the overalgebra in the restriction of a degenerate principal series of the group $\mathrm{GL}_4(\mathbb{R})$ to $\mathrm{GL}_2(\mathbb{R})$. Notice that canonical overgroups exist for all 10 series of real classical groups[4]. Moreover overgroups exist for all 52 series of classical semisimple symmetric spaces G/H, see [9], [15], see also [18], Addendum D.6. So the problem makes sense for all classical symmetric spaces.

[3] The problem of a decomposition of a tensor product $\rho_1 \otimes \rho_2$ of representations of a group G is a special case of a decomposition of restrictions. Indeed, $\rho_1 \otimes \rho_2$ is a representation of the group $G \times G$, and we must restrict this representation to the diagonal G.

[4] Emphasize that in the present paper we consider groups GL and not SL, also we must consider groups $\mathrm{U}(p,q)$ and not $\mathrm{SU}(p,q)$.

Shturm–Liouville problems for difference operators in $L^2(\mathbb{R})$ in the imaginary direction

$$\lambda f(s) = a(s)f(s+i) + b(s)f(s) + c(s)f(s-i)$$

arise in a natural way in the theory of hypergeometric orthogonal polynomials, see, e.g., [1], [8], apparently a first example (the Meixner–Pollaszek system) was discovered by J. Meixner in 1930s. On such operators with continuous spectra see [16], [6], [19]. See also a multi-rank work of I. Cherednik [2] on Harish-Chandra spherical transforms.

1.6. The Fourier transform on $\mathrm{GL}_2(\mathbb{C})$. For a detailed exposition of representations of the Lorentz group, see [14]. For ν, $\nu' \in \mathbb{C}$ satisfying $\nu - \nu' \in \mathbb{Z}$ we define the function $z^{\nu\|\nu'}$ on the multiplicative group of $\mathbb{C}$ by

$$z^{\nu\|\nu'} := z^{\nu}z^{\nu'} := |z|^{2\nu}\,\overline{z}^{\nu'-\nu}.$$

Denote by Δ the set of all $(\mu_1, \mu_1'; \mu_2, \mu_2') \in \mathbb{C}^4$ such that $\mu_1 - \mu_1' \in \mathbb{Z}$, $\mu_2 - \mu_2' \in \mathbb{Z}$. For $(\mu_1, \mu_1'; \mu_2, \mu_2') \in \Delta$ we define a representation of $\mathrm{GL}_2(\mathbb{C})$ in the space of functions on $\mathbb{C}$ by

$$T_{\mu_1,\mu_1';\mu_2,\mu_2'} \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} \varphi(t) =$$

$$= \varphi\Big(\frac{x_{12}+tx_{22}}{x_{11}+tx_{21}}\Big)(x_{11}+tx_{21})^{-1+\mu_1-\mu_2\|-1+\mu_1'-\mu_2'} \det \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}^{1/2+\mu_2\|1/2+\mu_2'}.$$

We consider the space of C^∞-smooth functions φ on $\mathbb{C}$ such that

$$\varphi(-t^{-1})t^{-1+\mu_1-\mu_2\|-1+\mu_1'-\mu_2'}$$

is C^∞-smooth at 0. These representations form the (nonunitary) *principal series*.

It is convenient to complexify the Lie algebra of $\mathrm{GL}_2(\mathbb{C})$,

$$\mathfrak{gl}_2(\mathbb{C})_{\mathbb{C}} \simeq \mathfrak{gl}_2(\mathbb{C}) \oplus \mathfrak{gl}_2(\mathbb{C}).$$

Under this isomorphism, the operators of the Lie algebra act in our representation by

$$L_{11} = -t\frac{d}{dt} + (-1/2 + \mu_1), \qquad L_{12} = \frac{d}{dt}, \tag{1.8}$$

$$L_{21} = -t^2\frac{d}{dt} + t(-1 + \mu_1 - \mu_2), \qquad L_{22} = t\frac{d}{dt} + (1/2 + \mu_2), \tag{1.9}$$

$$\overline{L}_{11} = -\overline{t}\frac{d}{d\overline{t}} + (-1/2 + \mu_1'), \qquad \overline{L}_{12} = \frac{d}{d\overline{t}}, \tag{1.10}$$

$$\overline{L}_{21} = -\overline{t}^2\frac{d}{d\overline{t}} + \overline{t}(-1 + \mu_1' - \mu_2'), \qquad \overline{L}_{22} = \overline{t}\frac{d}{d\overline{t}} + (1/2 + \mu_2'). \tag{1.11}$$

Formally, we have duplicated expressions (1.2)–(1.3).

If

$$\operatorname{Re}(\mu_1+\mu_1')=0,\qquad \operatorname{Re}(\mu_2+\mu_2')=0, \tag{1.12}$$

then the representation $T_{\mu_1,\mu_1';\mu_2,\mu_2'}$ is unitary in L^2. Denote by Δ_{tempered} the set of such tuples $(\mu_1,\mu_1';\mu_2,\mu_2')$, it is a union of a countable family of parallel 2-dimensional real planes in $\mathbb{C}^4$, we equip it by a natural Lebesgue measure $d\lambda(\mu)$.

For any compactly supported smooth function F on $\mathrm{GL}_2(\mathbb{C})$ we define its *Fourier transform* as an operator-valued function on Δ given by

$$T_{\mu_1,\mu_1';\mu_2,\mu_2'}(F)=\int_{\mathrm{GL}_2(\mathbb{C})} F(X)\,|\det X|^{-4}\prod_{k,l=1,2} d\operatorname{Re}x_{kl}\,d\operatorname{Im}x_{kl}.$$

The Plancherel formula is the following identity

$$\langle F_1,F_2\rangle_{L^2(\mathrm{GL}_2(\mathbb{C}))}=-C\cdot\int_{\Delta_{\text{tempered}}}\operatorname{tr}\big(T_{\mu_1,\mu_1';\mu_2,\mu_2'}(F_1)T_{\mu_1,\mu_1';\mu_2,\mu_2'}(F_2)\big)^*\times \\ \times(\mu_1-\mu_2)(\mu_1'-\mu_2')d\lambda(\mu),$$

where C is an explicit constant.

Denote by $K(t,s|\mu_1,\mu_1';\mu_2,\mu_2')$ the kernel of the operator $T_{\mu_1,\mu_1';\mu_2,\mu_2'}(F)$,

$$T_{\mu_1,\mu_1';\mu_2,\mu_2'}(F)\varphi(t)=\int_{\mathbb{C}} K(t,s|\mu_1,\mu_1';\mu_2,\mu_2')\varphi(s)\,ds.$$

1.7. Overgroup for $\mathrm{GL}_2(\mathbb{C})$. Consider the complex Grassmannian $\mathrm{Gr}_{4,2}(\mathbb{C})$ of 2-dimensional planes in $\mathbb{C}^4$, again the set $\mathrm{Mat}_2(\mathbb{C})$ is an open dense set on $\mathrm{Gr}_{4,2}(\mathbb{C})$. For σ, $\sigma'\in i\mathbb{R}$ consider a unitary representation $R_{\sigma,\sigma'}$ of $\mathrm{GL}_4(\mathbb{C})$ in $L^2\big(\mathrm{Mat}_2(\mathbb{C})\big)$ given by

$$R_{\sigma,\sigma'}F(X)=\begin{pmatrix}A&B\\C&D\end{pmatrix}F(X)=F\big((A+XC)^{-1}(B+XD)\big)\times \\ \times\ |\det(A+XC)|^{-2+2\sigma\|-2+2\sigma'}\left|\det\begin{pmatrix}A&B\\C&D\end{pmatrix}\right|^{1-\sigma\|1-\sigma'}.$$

We define a unitary operator $J: L^2\big(\mathrm{Mat}_2(\mathbb{C})\big)\to L^2\big(\mathrm{Mat}_2(\mathbb{C})\big)$ by

$$J_{\sigma,\sigma'}F(X)=F(X)(\det X)^{-1+\sigma\|-1+\sigma'}.$$

In this way we get a unitary representation $U_{\sigma,\sigma'}=J_{\sigma,\sigma'}R_{\sigma,\sigma'}J_{\sigma,\sigma'}^{-1}$ of $\mathrm{GL}_4(\mathbb{C})$ in $L^2\big(\mathrm{Mat}_2(\mathbb{C})\big)$:

$$U_{\sigma\|\sigma'}\begin{pmatrix}A&B\\C&D\end{pmatrix}F(X)=F\big((A+XC)^{-1}(B+XD)\big)\times \\ \times\left|\frac{\det(A+XC)\det(B+XD)}{\det X\det\begin{pmatrix}A&B\\C&D\end{pmatrix}}\right|^{-1+\sigma\|-1+\sigma'}.$$

1.8. Formulas for $\mathrm{GL}_2(\mathbb{C})$. We wish to write the action of the Lie algebra

$$\mathfrak{gl}_4(\mathbb{C})_{\mathbb{C}} \simeq \mathfrak{gl}_4(\mathbb{C}) \oplus \mathfrak{gl}_4(\mathbb{C})$$

in the Plancherel decomposition of $\mathrm{GL}_2(\mathbb{C})$. Denote the standard generators of $\mathfrak{gl}_4(\mathbb{C}) \oplus 0$ and $0 \oplus \mathfrak{gl}_4(\mathbb{C})$ by E_{kl} and $\overline{E}_{kl}$ respectively. Define the following shift operators

$$V_1 K(t,s;\mu_1,\mu_1',\mu_2,\mu_2') = K(t,s;\mu_1+1,\mu_1',\mu_2,\mu_2');$$
$$V_1' K(t,s;\mu_1,\mu_1',\mu_2,\mu_2') = K(t,s;\mu_1,\mu_1'+1,\mu_2,\mu_2'),$$

and similar operators V_2 and V_2' shifting μ_2 and μ_2'.

Then

$$E_{14} = \frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\,V_1^{-1} - \frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\,V_2^{-1};$$
$$\overline{E}_{14} = \frac{-1/2-\sigma'+\mu_1'}{\mu_1'-\mu_2'}\,\frac{\partial}{\partial \overline{s}}\,(V_1')^{-1} - \frac{-1/2-\sigma'+\mu_2'}{\mu_1'-\mu_2'}\,\frac{\partial}{\partial \overline{t}}\,(V_2')^{-1};$$
$$E_{32} = \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\,V_1 + \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\,V_2;$$
$$\overline{E}_{32} = \frac{1/2+\mu_1'+\sigma'}{\mu_1'-\mu_2'}\,\frac{\partial}{\partial \overline{t}}\,V_1' + \frac{1/2+\mu_2'+\sigma'}{\mu_1'-\mu_2'}\,\frac{\partial}{\partial \overline{s}}\,V_2'.$$

2 Calculations

2.1. The expression for kernel.

Lemma 2.1 *The kernel $K(\cdot)$ of an integral operator $T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}(F)$ is given by the formula*

$$K(t,s|\mu_1,\varepsilon_1;\mu_2,\varepsilon_2) =$$
$$= \iiint\limits_{\mathbb{R}^3} F\big(u-tv, su-stv-tw, v, sv+w\big)\, u^{-3/2+\mu_1/\!/\varepsilon_1}\, w^{-3/2+\mu_2/\!/\varepsilon_2}\, du\, dv\, dw. \tag{2.1}$$

For $F \in C_0^\infty\big(\mathrm{GL}_2(\mathbb{R})\big)$ the integration is actually taken over a bounded domain.

The integral converges if $\operatorname{Re}\mu_1 > 1/2$, $\operatorname{Re}\mu_2 > 1/2$. For fixed ε_1, ε_2 it has a meromorphic continuation to the whole complex plane (μ_1,μ_2) with poles on the hyperplanes $\mu_1 = -1/2-k$, $\mu_2 = -1/2-k$, where $k = 0, 1, 2, \ldots$ (see, e.g., [4], §I.3).

PROOF. By the definition

$$\int\limits_{\mathbb{R}} T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}(F)\varphi(t)\ \psi(t)\,dt =$$

$$= \int\limits_{\mathbb{R}} \int\limits_{\mathrm{Mat}_2(\mathbb{R})} F(x_{11},x_{12},x_{21},x_{22})\varphi\Big(\frac{x_{12}+tx_{22}}{x_{21}+tx_{12}}\Big)(x_{11}+tx_{21})^{-1+\mu_1-\mu_2/\!\!/\varepsilon_1-\varepsilon_2}\times$$

$$\times\,(x_{11}x_{22}-x_{12}x_{21})^{1/2+\mu_2/\!\!/\varepsilon_2}\,\frac{dx_{11}\,dx_{12}\,dx_{21}\,dx_{22}}{(x_{11}x_{22}-x_{12}x_{21})^2}\,dt.$$

In the interior integral, we pass from the variables x_{11}, x_{12}, x_{21}, x_{22} to new variables u, v, w, s defined by

$$\begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} = \begin{pmatrix} 1 & -t \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u & 0 \\ v & w \end{pmatrix} \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} \tag{2.2}$$

or

$$x_{11} = u - tv, \qquad x_{12} = su - stv - tw, \qquad x_{21} = v, \qquad x_{22} = sv + w.$$

The Jacobi matrix of this transformation is triangular, and the Jacobian is $|u|$. The inverse transformation is

$$u = x_{11} + tx_{21}, \quad v = x_{21}, \quad w = \frac{x_{11}x_{22} - x_{12}x_{21}}{x_{11} + tx_{21}}, \qquad s = \frac{x_{12} + tx_{22}}{x_{11} + tx_{21}}.$$

We also have $x_{11}x_{22} - x_{12}x_{21} = uw$.

After the change of variables we come to

$$\iint_{\mathbb{R}^2} K(t,s|\mu_1,\varepsilon_1;\mu_2,\varepsilon_2)\,\varphi(s)\,\psi(t)\,ds\,dt,$$

where $K(\cdot)$ is given by (2.1).

A function F has a compact support in $\mathbb{R}^4 \setminus \{x_{11}x_{22} - x_{12}x_{21} = 0\}$. So, actually, $x_{21} = v$, $x_{11} = u - tv$, $x_{22} = w + sv$ are contained in a bounded domain. This implies the second claim of the lemma. □

2.2. Preliminary remarks. Below F denotes

$$F := F(u - tv, su - stv - tw, v, sv + w).$$

Also, $\partial_{11}F$, $\partial_{12}F$, etc. denote

$$\partial_{11}F := \frac{\partial}{\partial x_{11}} F(x_{11},x_{12},x_{21},x_{22})\Big|_{\substack{x_{11}=u-tv \quad x_{12}=su-stv-tw,\\ x_{21}=v, \qquad\quad x_{22}=sv+w}}$$

etc. Partial derivatives of F are

$$\frac{\partial}{\partial u}F = \partial_{11}F + s\,\partial_{12}F; \tag{2.3}$$

$$\frac{\partial}{\partial v}F = -t\,\partial_{11}F - st\,\partial_{12}F + \partial_{21}F + s\,\partial_{22}F; \tag{2.4}$$

$$\frac{\partial}{\partial w}F = -t\,\partial_{12}F + \partial_{22}F, \tag{2.5}$$

and

$$\frac{\partial}{\partial s}F = (u - vt)\,\partial_{12}F + v\,\partial_{22}F; \tag{2.6}$$

$$\frac{\partial}{\partial t}F = -v\,\partial_{11}F - (w + sv)\,\partial_{12}F. \tag{2.7}$$

Also, notice that

$$\Big(y^{\nu/\!\!/\delta}\Big)' = \nu y^{\nu-1/\!\!/\delta+1}.$$

2.3. A verification of the formula for E_{14}. It is easy to verify that the operator e_{14} in $C_0^\infty\big(\mathrm{GL}_2(\mathbb{R})\big)$ is given by

$$e_{14} = \frac{\partial}{\partial x_{12}} - (-1+\sigma)\frac{x_{21}}{x_{11}x_{22} - x_{12}x_{21}}.$$

Therefore,

$$\begin{aligned} T_{\mu_1,\varepsilon_1;\mu_2,\varepsilon_2}(e_{14}F) = \\ = \iiint \partial_{12}F\; u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw - \\ - (-1+\sigma)\iiint F\frac{v}{uw}\,u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw, \end{aligned} \tag{2.8}$$

(the integration is taken over $\mathbb{R}^3$ on default).

We must verify that (2.8) coincides with

$$E_{14}K = \Big(\frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\,V_1^- + \frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\,V_2^-\Big)K \tag{2.9}$$

Below we establish two formulas

$$\begin{aligned} \iiint \partial_{12}F\; u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw \;-\; \frac{\partial}{\partial s}V_1^- K = \\ = (-3/2+\mu_2)\iiint F\;\frac{v}{uw}\,u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw, \end{aligned} \tag{2.10}$$

$$\begin{aligned} \iiint \partial_{12}F\; u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw \;+\; \frac{\partial}{\partial t}V_2^- K = \\ = (-3/2+\mu_1)\iiint F\;\frac{v}{uw}\,u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw. \end{aligned} \tag{2.11}$$

Considering the sum of (2.10) and (2.11) with coefficients $\frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2}$ and $\frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2}$ we get coincidence of (2.8) and (2.9); for this, we use he identities

$$\frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2} - \frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2} = 1;$$

$$(-3/2+\mu_2)\cdot\frac{-1/2-\sigma+\mu_1}{\mu_1-\mu_2} - (-3/2+\mu_1)\cdot\frac{-1/2-\sigma+\mu_2}{\mu_1-\mu_2} = 1-\sigma.$$

Now let us check (2.10). The following identity can be verified by a straightforward calculation (with (2.6) and (2.5)):

$$\partial_{12}F - \frac{1}{u}\frac{\partial}{\partial s}F = -\frac{v}{u}\frac{\partial}{\partial w}F.$$

Therefore the left-hand side of (2.10) equals to

$$\iiint \Big[-\frac{v}{u}\frac{\partial}{\partial w}F\Big] \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw.$$

We integrate this expression by parts in the variable w and come to (2.10).

To check (2.11), we verify the identity

$$\partial_{12}F + \frac{1}{v}\frac{\partial}{\partial t}F = \frac{v}{w}\frac{\partial}{\partial u}F$$

and after this integrate by parts as above.

2.4. A verification of the formula for E_{32}. We have

$$e_{32} = -\Big(x_{11}x_{21}\frac{\partial}{\partial x_{11}} + x_{11}x_{22}\frac{\partial}{\partial x_{12}} + x_{21}^2\frac{\partial}{\partial x_{21}} + x_{21}x_{22}\frac{\partial}{\partial x_{22}}\Big) + (-1+\sigma)x_{21}.$$

Therefore,

$$T(e_{32}K) = \iiint \big(G + (-1+\sigma)vF\big) \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw, \quad (2.12)$$

where

$$G = (u-vt)v\,\partial_{11}F + (u-vt)(w+vs)\,\partial_{12}F + v^2\,\partial_{21}F + v(w+vs)\,\partial_{22}F.$$

On the other hand,

$$E_{32}K = \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\iiint u\frac{\partial}{\partial t}F \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw + \\ + \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\iiint w\frac{\partial}{\partial s}F \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw. \quad (2.13)$$

We must verify that (2.12) and (2.12) are equal. As in the previous subsection, this statement is reduced to a pair of identities

$$\iiint \big(G + (-1+\sigma)vF - u\frac{\partial}{\partial t}F\big) \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw = \\ = (1/2+\mu_2+\sigma)\iiint vF \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw; \quad (2.14)$$

$$\iiint \big(G + (-1+\sigma)vF + w\frac{\partial}{\partial s}F\big) \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw = \\ = (1/2+\mu_1+\sigma)\iiint vF \cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1} w^{-3/2+\mu_2/\!\!/\varepsilon_2}\, du\, dv\, dw. \quad (2.15)$$

Let us verify (2.14). It can be easily checked (with (2.7), (2.4), (2.5)) that

$$G - u\frac{\partial}{\partial t}F = -v^2\frac{\partial}{\partial v}F - vw\frac{\partial}{\partial w}F.$$

We substitute this to the left-hand side of (2.14) and come to

$$\iiint \bigl(-v^2\frac{\partial}{\partial v}F - vw\frac{\partial}{\partial w}F + (-1+\sigma)vF\bigr)\times \\ \times\ u^{-3/2+\mu_1/\!\!/\varepsilon_1}w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw. \quad (2.16)$$

Integrating by parts, we get

$$\iiint F\cdot\frac{\partial}{\partial v}\Bigl(v^2\Bigr)\cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1}w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw+ \\ + \iiint vF\cdot u^{-3/2+\mu_1/\!\!/\varepsilon_1}\frac{\partial}{\partial v}\Bigl(w^{-1/2+\mu_2/\!\!/\varepsilon_2+1}\Bigr)\,du\,dv\,dw+ \\ + (-1+\sigma)\iiint vF\cdot\ u^{-3/2+\mu_1/\!\!/\varepsilon_1}w^{-3/2+\mu_2/\!\!/\varepsilon_2}\,du\,dv\,dw.$$

After a summation we come to the right-hand side of (2.14).

A proof of (2.15) is similar, we use the identity

$$G + v\frac{\partial}{\partial s}F = -v^2\frac{\partial}{\partial v}F - -uv\frac{\partial}{\partial u}F$$

and repeat the same steps.

2.5. Table of formulas. First, we present formulas for the action of the Lie algebra $\mathfrak{gl}_4$ corresponding to U_σ, see (1.6). Denote generators of $\mathfrak{gl}_4$ by e_{kl}, where $1\leqslant k,l\leqslant 4$. Denote by ∂_{pq} the partial derivatives $\frac{\partial}{\partial x_{pq}}$, where $p,q=1,2$. The generators e_{kl} naturally split into 4 groups corresponding to blocks A, B, C, D in (1.6).

a) Generators corresponding to the block A form a Lie algebra $\mathfrak{gl}_2$:

$$e_{11} = -x_{11}\partial_{11} - x_{12}\partial_{12}, \qquad e_{12} = -x_{21}\partial_{11} - x_{22}\partial_{12},$$
$$e_{21} = -x_{11}\partial_{21} - x_{12}\partial_{22}, \qquad e_{22} = -x_{21}\partial_{21} - x_{22}\partial_{22}.$$

b) Generators corresponding to the block D also form a Lie algebra $\mathfrak{gl}_2$:

$$e_{33} = x_{11}\partial_{11} + x_{21}\partial_{21}, \qquad e_{34} = x_{11}\partial_{12} + x_{21}\partial_{22},$$
$$e_{43} = x_{12}\partial_{11} + x_{22}\partial_{21}, \qquad e_{44} = x_{12}\partial_{12} + x_{22}\partial_{22}.$$

c) Elements corresponding to the block B form a 4-dimensional Abelian Lie algebra:

$$e_{13} = \partial_{11} + (-1+\sigma)\frac{x_{22}}{\det X}, \qquad e_{14} = \partial_{12} - (-1+\sigma)\frac{x_{21}}{\det X},$$
$$e_{23} = \partial_{21} - (-1+\sigma)\frac{x_{12}}{\det X}, \qquad e_{24} = \partial_{22} + (-1+\sigma)\frac{x_{11}}{\det X}.$$

c) Elements corresponding to the block C also form a 4-dimensional Abelian Lie algebra:

$$
\begin{aligned}
e_{31} &= -(x_{11}^2\partial_{11} + x_{11}x_{12}\partial_{12} + x_{11}x_{21}\partial_{21} + x_{12}x_{21}\partial_{22}) + (-1+\sigma)x_{11},\\
e_{32} &= -(x_{11}x_{21}\partial_{11} + x_{11}x_{22}\partial_{12} + x_{21}^2\partial_{21} + x_{21}x_{22}\partial_{22}) + (-1+\sigma)x_{21},\\
e_{41} &= -(x_{11}x_{12}\partial_{11} + x_{12}^2\partial_{12} + x_{11}x_{22}\partial_{21} + x_{12}x_{22}\partial_{22}) + (-1+\sigma)x_{12},\\
e_{42} &= -(x_{12}x_{21}\partial_{11} + x_{12}x_{22}\partial_{12} + x_{21}x_{22}\partial_{21} + x_{22}^2\partial_{22}) + (-1+\sigma)x_{22}.
\end{aligned}
$$

Denote by E_{kl} the corresponding operators E_{kl} on kernels K. Formulas for operators of groups a), b) immediately follow from the definition of the Fourier transform,

$$
\begin{aligned}
E_{11} &= -t\frac{\partial}{\partial t} - (1/2-\mu_1), & E_{12} &= \frac{\partial}{\partial t},\\
E_{21} &= -t^2\frac{\partial}{\partial t} + (-1+\mu_1-\mu_2)t, & E_{22} &= t\frac{\partial}{\partial t} + (1/2+\mu_2),
\end{aligned}
$$

and

$$
\begin{aligned}
E_{33} &= -s\frac{\partial}{\partial s} - (1/2+\mu_1), & E_{34} &= \frac{\partial}{\partial s},\\
E_{43} &= -s^2\frac{\partial}{\partial s} + (-1-\mu_1+\mu_2)s, & E_{44} &= s\frac{\partial}{\partial s} + (1/2-\mu_2).
\end{aligned}
$$

Next,

$$
\begin{aligned}
E_{13} &:= \frac{1/2-\mu_2+\sigma}{\mu_1-\mu_2}\, s\frac{\partial}{\partial t}\, V_2^- + \frac{1/2-\mu_1+\sigma}{\mu_1-\mu_2}\Big(\mu_1-\mu_2+s\frac{\partial}{\partial s}\Big)\, V_1^-,\\
E_{14} &= -\frac{1/2+\sigma-\mu_1}{\mu_1-\mu_2}\,\frac{\partial}{\partial s}\, V_1^- - \frac{1/2+\sigma-\mu_2}{\mu_1-\mu_2}\,\frac{\partial}{\partial t}\, V_2^-,\\
E_{23} &:= \frac{1/2-\mu_2+\sigma}{\mu_1-\mu_2}\, s\left(-(\mu_1-\mu_2)+t\frac{\partial}{\partial t}\right) V_2^- +\\
&\qquad\qquad \frac{1/2-\mu_1+\sigma}{\mu_1-\mu_2}\, t\left(\mu_1-\mu_2+s\frac{\partial}{\partial s}\right) V_1^-,\\
E_{24} &= -\frac{1/2+\sigma-\mu_1}{\mu_1-\mu_2} t\,\frac{\partial}{\partial s}\, V_1^- + \frac{1/2+\mu_2-\sigma}{\mu_1-\mu_2}\left(\mu_1-\mu_2+t\frac{\partial}{\partial s}\right) V_2^-.
\end{aligned}
$$

and

$$
\begin{aligned}
E_{31} &:= \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\Big(\mu_1-\mu_2-t\frac{\partial}{\partial t}\Big)\, V_1^+ - \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\, t\frac{\partial}{\partial s}\, V_2^+,\\
E_{32} &= \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\frac{\partial}{\partial t}\, V_1^+ + \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\frac{\partial}{\partial s}\, V_2^+,\\
E_{41} &= \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}\, s\Big(\mu_1-\mu_2-t\frac{\partial}{\partial t}\Big)V_1^+ - \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\, t\left(\mu_1-\mu_2+s\frac{\partial}{\partial s}\right)V_2^+,\\
E_{42} &= \frac{1/2+\mu_1+\sigma}{\mu_1-\mu_2}s\frac{\partial}{\partial t}V_1^+ + \frac{1/2+\mu_2+\sigma}{\mu_1-\mu_2}\Big(\mu_1-\mu_2+s\frac{\partial}{\partial s}\Big)V_2^+.
\end{aligned}
$$

2.6. The case of $\mathrm{GL}_2(\mathbb{C})$. Notice that formulas in Subsections 1.1–1.4 for $\mathrm{SL}_2(\mathbb{R})$ and in Subsections 1.6–1.8 are very similar, except the Plancherel formulas.

The analog of formula (2.1) is

$$K(t,s|\mu_1,\mu_1';\mu_2,\mu_2') =$$
$$= \iiint\limits_{\mathbb{C}^3} F\big(u-tv, su-stv-tw, v, sv+w\big)\, u^{-3/2+\mu_1\|-3/2+\mu_1'} w^{-3/2+\mu_2\|-3/2+\mu_2'} \times$$
$$\times\ d\operatorname{Re} u\, d\operatorname{Im} u\, d\operatorname{Re} v\, d\operatorname{Im} v\, d\operatorname{Re} w\, d\operatorname{Im} w. \quad (2.17)$$

Its derivation is based on the same change of variables (2.2), its real Jacobian is $u\overline{u}$. A further calculation one-to-one follows the calculation for $\mathrm{GL}_2(\mathbb{R})$.

References

[1] Andrews G. R., Askey R., Roy R., *Special functions*, Cambridge Univ. Press, Cambridge, 1999

[2] Cherednik I., *Inverse Harish-Chandra transform and difference operators.* Int. Math. Res. Not. 1997, No.15, 733-750 (1997).

[3]

[4] Gelfand, I. M.; Shilov, G. E. *Generalized functions. Vol. 1: Properties and operations.* Providence, RI: AMS Chelsea Publishing (2016).

[5] Gelfand, I. M.; Graev, M. I.; Vilenkin, N. Ya. *Generalized functions. Vol. 5: Integral geometry and representation theory.* Providence, RI: AMS Chelsea Publishing, 2016.

[6] Groenevelt, W. *The Wilson function transform.* Int. Math. Res. Not. 2003, no. 52, 2779-2817.

[7] Howe, R.; Tan, Eng Chye *Non-Abelian harmonic analysis. Applications of* $\mathrm{SL}(2,\mathbb{R})$. Springer-Verlag, (1992).

[8] Koekoek R., Swarttouw R. F. *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue.* Delft University of Technology Faculty of Information Technology and Systems. Department of Technical Mathematics and Informatics Report no. 98-17 1998. Available via `http://homepage.tudelft.nl/11r49/askey.html`

[9] Makarevich, B. O. *Open symmetric orbits of reductive groups in symmetric R-spaces*, Math. USSR-Sb., 20:3 (1973), 406-418

[10] Molchanov, V. F. *Canonical representations and overgroups for hyperboloids of one sheet and Lobachevsky spaces.* Acta Appl. Math. 86 (2005), no. 1-2, 115-129.

[11] Molchanov, V. F. *Canonical representations and overgroups for hyperboloids.* Funct. Anal. Appl. 39 (2005), no. 4, 284-295

[12] Molchanov, V. F. *Canonical representations on Lobachevsky spaces: an interaction with an overalgebra.* Acta Appl. Math. 99 (2007), no. 3, 321-337.

[13] Molchanov, V. F. *Fourier and Poisson transforms for tensor products.* Functional Analysis and Its Applications, 2015, 49:4, 279-288.

[14] Naimark M. A. *Linear representations of Lorentz group.* Pergamon Press, 1964.

[15] Neretin, Yu. A. *Pseudo-Riemannian symmetric spaces: uniform realizations and open embeddings into Grassmannians,* Zap. Nauchn. Sem. POMI, 256, POMI, St. Petersburg, 1999, 145-167; J. of Math. Sciences (New York), 2001, 107:5, 4248-4264.

[16] Neretin, Yu. A. *Index hypergeometric transform and imitation of analysis of Berezin kernels on hyperbolic spaces,* Sbornik Math., 192:3 (2001), 403–432

[17] Neretin, Yu. A. *The action of an overalgebra in the Plancherel decomposition and shift operators in an imaginary direction.* Izv. Math. 66 (2002), no. 5, 1035-1046.

[18] Neretin, Yu. A. *Lectures on Gaussian integral operators and classical groups.* European Mathematical Society (EMS), Zürich, 2011.

[19] Neretin, Yu. A. *Difference Sturm-Liouville problems in the imaginary direction.* J. Spectr. Theory 3 (2013), no. 3, 237-269.

[20] Neretin, Yu. A. *Restriction of representations of* $\mathrm{GL}(n+1,\mathbb{C})$ *to* $\mathrm{GL}(n,\mathbb{C})$ *and action of the Lie overalgebra.* Preprint, arXiv:1510.03611

Math. Dept., University of Vienna;
Institute for Theoretical and Experimental Physics (Moscow);
MechMath Dept., Moscow State University;
Institute for Information Transmission Problems.
URL: http://mat.univie.ac.at/∼neretin/